

Erratum

In the original article by Lu, et al., entitled “Stimulating Effect of a Newly Synthesized Sulfonamido-Based gallate on Articular Chondrocytes *in Vitro*” [Cell Physiol Biochem 2015;37(3):1196-1209, DOI: 10.1159/000430243], there are some errors in Fig. 3 and Fig. 5. The images of cell viability in Fig. 3A: TGF- β 1 at day 2 and 1.06×10^{-7} M at day 6, and in Fig. 5: phalloidin/Hoechst 33258 staining 0M, 1.06×10^{-8} M, 1.06×10^{-7} M at day 2, 1.06×10^{-7} M and 1.06×10^{-6} M at day 4, and 1.06×10^{-7} M at day 6 were mistakenly submitted. The correct Fig. 3 and 5 are displayed below.

The authors confirm that all of the results and conclusions of the article remain unchanged, as well as the figure legend.
The authors sincerely apologize for this mistake.

Fig. 3. (A) Confocal laser scanning microscopy images showing the viability of chondrocytes cultured *in vitro* alone (Control) or with ZXHA-C (1.06×10^{-8} M, 1.06×10^{-7} M, 1.06×10^{-6} M) and TGF- β 1 (T=15 ng/mL) for 2, 4 and 6 days. Cell seeding density: 2×10^4 /mL (original magnification $\times 40$, scale bar was 100 μ m). (B) Cell number calculated and analyzed on the basis of FDA/PI staining images. Cell numbers were counted in one image by using the “cell calculating” toolbar in the Nikon A1 software. The data represent the mean \pm SD of three independent ulcure experiments. Bars with different letters are significantly different from each other at $P < 0.05$.




Fig. 5. Phalloidin/Hoechst 33258 staining images showing the morphology of chondrocytes cultured *in vitro* alone (Control) or with ZXHA-C (1.06×10^{-8} M, 1.06×10^{-7} M, 1.06×10^{-6} M) and TGF- β 1 (T=15 ng/mL) for 2, 4 and 6 days. Cell seeding density: 2×10^4 /mL (Original magnification $\times 100$, scale bar was 100 μ m).

