


Supplementary Material

Osteoprotegerin Expression in Liver is Induced by IL13 through TGF β

Adhyatmika Adhyatmika^{a,b} Kurnia S. S. Putri^{c,d} Emilia Gore^c Keri A. Mangnus^a
Catharina Reker-Smit^a Detlef Schuppan^{e,f} Leonie Beljaars^c Peter Olinga^c
Barbro N. Melgert^{g,h}

^aDepartment of Pharmacokinetics, Toxicology, and Targeting, Groningen Research Institute of Pharmacy, University of Groningen, Groningen, The Netherlands, ^bDepartment of Pharmaceutics, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia, ^cDepartment of Pharmaceutical Technology and Biopharmacy, Groningen Research Institute of Pharmacy, University of Groningen, Groningen, The Netherlands, ^dFaculty of Pharmacy, Universitas Indonesia, Depok, West Java, Indonesia, ^eInstitute of Translational Immunology, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany, ^fDivision of Gastroenterology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA, ^gDepartment of Molecular Pharmacology, Groningen Research Institute of Pharmacy, University of Groningen, Groningen, The Netherlands, ^hGroningen Research Institute for Asthma and COPD, University Medical Center Groningen, Groningen, The Netherlands



Supplementary Figure S1. Treatments did not compromise viability mouse liver slices. Treatment of liver slices for 48 hours with compounds in our experiments did not significantly compromise the viability of the mouse liver slices used in our study (n=6). Treated groups were compared with the control using a Kruskal-Wallis test corrected for multiple testing, $p < 0.05$ was considered significant.