Cellular Physiology	Cell Physiol Biochem 2022;56:310-311	
and Biochemistry	DOI: 10.33594/000000536	© 2022 The Author(s) Published by Cell Physiol Biochem Press GmbH&Co. KG, Duesseldorf www.cellphysiolbiochem.com

Erratum

In the article "Stimulating Effect of a Newly Synthesized Sulfonamido-Basedgallate on Articular Chondrocytes *in Vitro*" [Cell Physiol Biochem 2015;37:1196-1209; DOI: 10.1159/000430243] by Lu et al., incorrect representative images were included in Figure 3A (1.06x10-8 M at day 2, 0 M at day 4, 1.06x10-8 M at day 4, 1.06x10-7 M at day 4 and 1.06x10-8 M at day 6) and Figure 4 (1.06x10-7 M at day 4). This was as a result of poor record keeping at the time of the experiment. The correct files have been verified by review of the laboratory notebooks and the authors state that the results and conclusions of the article remain unchanged. The authors apologize for any inconvenience caused.

The corrected Figure 3 and Figure 4 are shown here. The corrected Figure 3 includes images corrected in a previous Erratum (https://doi.org/10.33594/000000137).

Fig. 3. (A) Confocal laser scanning microscopy images showing the viability of chondrocytes cultured *in vitro* alone (Control) or with ZXHA-C $(1.06 \times 10^{-8} \text{ M})$ 1.06×10^{-7} M, 1.06×10^{-6} M) and TGF- β 1 (T=15 ng/mL) for 2, 4 and 6 days. Cell seeding density: 2×10⁴/mL (original magnification \times 40, scale bar was 100 μ m). (B) Cell number calculated and analyzed on the basis of FDA/ PI staning images. Cell numbers were counted in one image by using the "cell calculating" toolbar in the Nikon A1 software. The data represent the mean \pm SD of three independent ulcutre experiments. Bars with different letters are significantly different from each other at P < 0.05.



Cellular Physiology	Cell Physiol Biochem 2022;56:310-311		
and Biochemistry	DOI: 10.33594/000000536	© 2022 The Author(s). Published by Cell Physiol Biochem Press GmbH&Co. KG	311
······································	Erratum	•	-

Fig. 4. Hematoxylin-eosin staining images showing the morphology of chondrocytes cultured *in vitro* alone (Control) or with ZHA-C (1.05×10^{-8} M, 1.06×10^{-7} M, 1.06×10^{-6} M) and TGF- β 1 (T=15 ng/mL) for 2, 4 and 6 days. Cell seeding density: 2×10^{4} /mL (original magnification × 200, scale bar was 100 µm).

2d	Control	1.06×10-8 M	1.06×10-7 M	1.06×10-6 M	TGF-β1
4d					
6d					